# **RETRACTION NOTE**

**Open Access** 

# Retraction Note: Desflurane anesthesia worsens emergence agitation in adult patients undergoing thyroid surgery compared to sevoflurane anesthesia



Takeshi Suzuki<sup>1\*</sup>, Takuya Kurazumi<sup>1</sup>, Tomomi Ueda<sup>1</sup>, Hiromasa Nagata<sup>1</sup>, Takashige Yamada<sup>1</sup>, Shizuko Kosugi<sup>1</sup>, Saori Hashiguchi<sup>1</sup>, Koichi Ito<sup>2</sup> and Hiroshi Morisaki<sup>1</sup>

Correction to: JA Clin Rep (2017) 3:36 https://doi.org/10.1186/s40981-017-0106-5

## Retraction

The Editor-in-Chief has retracted this article [1]. The ethics committee approval was granted for an observational study and the need for patient consent was waived. However, the study design described is a randomized controlled trial and therefore patient consent should have been obtained. All authors agree with this retraction.

### **Author details**

<sup>1</sup>Department of Anesthesiology, Keio University School of Medicine, 35 Shinanomachi, Shinjuku-ku, Tokyo 160-8582, Japan. <sup>2</sup>Department of Surgery, Ito Hospital, Tokyo, Japan.

Published online: 14 February 2020

### Reference

 Suzuki T, Kurazumi T, Ueda T, Nagata H, Yamada T, Kosugi S, Hashiguchi S, Ito K, Morisaki H. Desflurane anesthesia worsens emergence agitation in adult patients undergoing thyroid surgery compared to sevoflurane anesthesia. JA Clin Rep. 2017;3:36. https://doi.org/10.1186/s40981-017-0106-5.

The original article can be found online at https://doi.org/10.1186/s40981-017-0106-5

<sup>&</sup>lt;sup>1</sup>Department of Anesthesiology, Keio University School of Medicine, 35 Shinanomachi, Shinjuku-ku, Tokyo 160-8582, Japan Full list of author information is available at the end of the article



<sup>\*</sup> Correspondence: takeshi-su@a7.keio.jp