CORRECTION

Open Access

Correction: The incidence of hypotension during general anesthesia: a single-center study at a university hospital

Nobuyuki Katori¹, Kentaro Yamakawa¹, Kotaro Kida¹, Yoshihiro Kimura¹, Shoko Fujioka¹ and Tsunehisa Tsubokawa^{1*}

Correction: JA Clin Rep 9, 23 (2023) https://doi.org/10.1186/s40981-023-00617-9

Following publication of the original article [1], the author reported that there are citations to Figs. 2 and 3 that should have not been included in the second paragraph of the 'Results' section. The paragraph should read: From

We also calculated the percentage of events that continued for more than or equal to 10 min at each hypotensive threshold using the data presented in Fig. 3. The percentage of hypotensive events that continued for more than or equal to 10 min at <75 mmHg, <65 mmHg, <55 mm Hg, and <45 mmHg was 81.4%, 57.6%, 18.4%, and 6.5%, respectively. Considering the incidence of IOH shown in Fig. 2, the incidence of IOH continuing for \geq 10 min in each hypotensive threshold described above was 78.9%, 49.7%, 8.9%, and 0.8%, respectively.

То

We also calculated the percentage of events that continued for more than or equal to 10 min at each hypotensive threshold: 81.4% for <75 mmHg, 57.6\% for <65 mmHg, 18.4% for 55 mmHg, and 6.5% for <45 mmHg,

The original article can be found online at https://doi.org/10.1186/s40981-023-00617-9.

*Correspondence: Tsunehisa Tsubokawa

¹ Department of Anesthesiology, The Jikei University School of Medicine,

2-25-8 Nishishinbashi, Minato-Ku, Tokyo 105-8461, Japan

Den Springer Open

© The Author(s) 2023. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit http://creativecommons.org/licenses/by/4.0/.

respectively. Hence, the incidence of IOH continuing for ≥ 10 min in each hypotensive threshold described above was 78.9%, 49.7%, 8.9%, and 0.8%, respectively.

The original article has been updated.

Published online: 01 June 2023

Reference

1. Katori N, Yamakawa K, Kida K, et al. The incidence of hypotension during general anesthesia: a single-center study at a university hospital. JA Clin Rep. 2023;9:23. https://doi.org/10.1186/s40981-023-00617-9.

⁻⁻⁻⁻⁻⁻

tsunetann@gmail.com